OK TO ENTER: /ESO/

Applicant : Nai-Kong CHEUNG Atty. Dkt. # : 639-C-PCT-US

USSN : 10/565,484 Art Unit : 1623

Filed : January 17, 2006 Date of Office Action: June 16, 2009
Examiner : Eric Olson Date of Response : September 9, 2009

Page : 3

AMENDMENT TO THE CLAIMS

This listing of claims will replace all prior listings or versions of claims in this application.

1-13. (Canceled)

- 14. (Previously Presented) A composition comprising:
 - (a) a composition comprising an antibody that binds to a cancer cell, and at least one pharmaceutically acceptable carrier, wherein the cancer is selected from the group consisting of neuroblastoma, melanoma, non-Hodgkin's lymphoma, breast cancer, Epstein-Barr related lymphoma, Hodgkin's lymphoma, and epidermoid carcinoma; and
 - (b) an orally administered composition comprising at least one pharmaceutically acceptable carrier and a soluble β -glucan in an amount effective to enhance the antitumor effect of said antibody, wherein the β -glucan comprises a β -1,3 backbone and at least one β -1,3 side chain of two or more glucose units linked to the backbone by a β -1,6 glycosidic bonds.
- 15. (Previously Presented) The composition of claim 14, wherein the β -glucan is isolated from yeast.
- 16. (Previously Presented) The composition of claim 14, wherein the β -glucan is isolated from Saccharomyces Cerevisiae.

Applicant : Nai-Kong CHEUNG Atty. Dkt. # : 639-C-PCT-US

USSN : 10/565,484 Art Unit : 1623

Filed : January 17, 2006 Date of Office Action: June 16, 2009
Examiner : Eric Olson Date of Response : September 9, 2009

Page: 4

17. (Currently Amended) The composition of claim 14, wherein the β -glucan has a molecular weight from about 10 kDa to about 350 kDa and is capable of inducing cytokines.

18. (Canceled)

- 19. (Previously Presented) The composition of claim 14, wherein the antibody is a monoclonal antibody or a complement-activating antibody.
- 20. (Previously Presented) The composition of claim 14, wherein the antibody binds to a cancer cell expressing an antigen selected from the group consisting of CD20, HER2, EGFR, GD2, and GD3.
- 21. (Previously Presented) The composition of claim 14, wherein the antibody is further capable of activating an antibody dependent cell-mediated cytotoxicity response.
- 22. (Previously Presented) A composition comprising:
 - (a) a composition comprising an antibody that binds to a cancer cell, and at least one pharmaceutically acceptable carrier, wherein the antibody binds to a cancer cell expressing an antigen selected from the group consisting of CD20, HER2, EGFR, GD2, and GD3; and
 - (b) an orally administered composition comprising at least one pharmaceutically acceptable carrier and a soluble β -glucan in an amount effective to enhance the antitumor effect of said antibody, wherein the β -glucan comprises a

Applicant : Nai-Kong CHEUNG Atty. Dkt. # : 639-C-PCT-US

USSN : 10/565,484 Art Unit : 1623

Filed : January 17, 2006 Date of Office Action: June 16, 2009
Examiner : Eric Olson Date of Response : September 9, 2009

Page : 5

 β -1,3 backbone and at least one β -1,3 side chain of two or more glucose units linked to the backbone by a β -1,6 glycosidic bond.

- 23. (Previously Presented) The composition of claim 22, wherein the β -glucan is isolated from yeast.
- 24. (Previously Presented) The composition of claim 22, wherein the β -glucan is isolated from Saccharomyces Cerevisiae.
- 25. (Currently Amended) The composition of claim 22, wherein the β -glucan has a molecular weight from about 10 kDa to about 350 kDa and is capable of inducing cytokines.
- 26. (Canceled)
- 27. (Previously Presented) The composition of claim 22, wherein the antibody is a monoclonal antibody or a complement-activating antibody.
- 28. (Previously Presented) The composition of claim 22, wherein the cancer cell is selected from the group consisting of neuroblastoma, melanoma, non-Hodgkin's lymphoma, breast cancer, Epstein-Barr related lymphoma, Hodgkin's lymphoma, and epidermoid carcinoma.
- 29. (Previously Presented) The composition of claim 22, wherein the antibody is further capable of activating an antibody dependent cell-mediated cytotoxicity response.